2016 MBL
Undergraduate Research Symposium

Thursday, August 18, 2016
Speck Auditorium
8:00 am to 5:30 pm

2016 MBL Undergraduate Research Symposium
August 18

10:20 am BREAK

Session Two±Chair: Lydia Mäthger

10:30 am Using Lipid Biomarkers to Understand Deep Ocean Organic Particle Flux
Leonard Shaw[1], Emily Maness[2], J.C. Weber[3], Maureen Conte[3]. *[1]University of Chicago, Chicago IL; [2]University of Tampa, Tampa FL; [3]Ecosystems Center, Marine Biological Laboratory, Woods Hole, MA*

10:45 am Decadal shifts in the vegetation structure Great Sippewissett Marsh in response to sea level rise.
Tynan Bowyer[1,2], Ivan Valiela[2], Javier Lloret[2], David Remsen[2], Simon Miner[2]. *[1]University of Chicago, Chicago, IL; [2]Marine Biological Lab, Woods Hole, MA.*

11:00 am Long term responses to high N loading in West Falmouth Harbor: the elusive role of eelgrass meadows in determining water quality.
Toby SantaMaria[1], Anne Giblin[2], and Melanie Hayn[2, 3]. *[1]Kenyon College Dept of Biology, Gambier OH 43022; [2]*

Dynamics of ***Asterias forbesi*** and ***A. vulgaris*** microbial communities in relation to sea star wasting disease (SSWD)

Assessing Microbial Metabolic Function and Circadian Rhythms Over Time And Space In Siders Pond

Petra K. Byl[1,2], Emily Reddington[2], Rika Anderson[3], Joseph J. Vallino[2], Julie A. Huber[2]
[1]University of Chicago, Chicago, IL; [2]Marine Biological Laboratory, Woods Hole, MA

Diverse bacterial and archaeal metabolic networks cycle life-essential elements throughout our biosphere and can locally influence the habitability of an ecosystem. However, we have limited knowledge on the dynamics that govern interactions between members of a microbial community in the environment over time and space. Here, we examined the structure and function of the microbial community in Siders Pond— a permanently stratified, meromictic kettle pond in Falmouth, MA— with a focus on the relationship between phototrophs and other members of the microbial community. We sampled 7 casts at 8 chemically and physically distinct depths over a 24-hour diel cycle and analyzed biogeochemical constituents as well as constructed metagenomic and metatranscriptomic libraries. Results show a vertical gradient of phototrophs over depth with mixotrophic and aerobic phototrophic lineages dominating the water column above 6 meters and anaerobic phototrophic lineages dominating the water column at 8 meters. Our metagenomic and geochemical data indicates that dissolved oxygen, microbial biomass, and regulatory circadian rhythm genes (kaiABC, ldpA, and cikA) peaked at 3-4 meters depth, where phototrophic cyanobacteria dominated the water column. In the lower water column, a coupled decrease in hydrogen sulfide and dissolved inorganic carbon during daylight hours provided evidence for a diel metabolic cycle in phototrophic green sulfur bacteria (*Chlorobi*). Based on our preliminary data, we hypothesize that microbial phototrophs express genes on an internal circadian rhythm. We are currently analyzing the metatranscriptome for the expression of regulatory circadian rhythm genes, phototrophic metabolism, and nutrient uptake. Combined with data from the spatial and temporal changes in water column geochemistry, results will be presented in the context of understanding the complex microbiome that drives biogeochemical cycling in Siders Pond.

Funding by NSF grant GEO-1451356

Inhibition of fast axonal transport in neurons is mediated by mutant huntingtin protein

[1,3]Karen G. Ebenezer, [2,3]Minsu Kang, [2,3]Gerardo Morfini, [2,3]Scott T. Brady
[1]CUNY Hunter College, New York City, NY; [2]University of Illinois at Chicago, Chicago, IL; and [3]Marine Biological Laboratory, Woods Hole, MA.

Huntington's disease (HD) is a fatal adult-onset neurodegenerative disease characterized by an involuntary jerky movement called Huntington's chorea as well as cognitive deficits and mood disorders. HD is caused by a mutation of a trinucleotide repeat in the huntingtin gene which results in an expansion of the polyglutamine region of the huntingtin protein (Htt). There is no known treatment for the disease, and it affects 1 in 10,000 people in the United States. A growing body of evidence indicates that the inhibition of fast axonal transport (FAT) by mutant Htt (mHtt) is a major pathogenic event in HD. Several studies using different model organisms of HD report inhibition of both anterograde and retrograde axonal transport of various molecular cargoes mediated by motor proteins kinesin and cytoplasmic dynein respectively. Using squid axoplasm isolated from squid giant axons, our laboratory demonstrated that mHtt leads to inhibition of FAT via activation of a mitogen activated protein kinase (MAPK) called c-Jun N-terminal kinase 3 (JNK3), which phosphorylates kinesin and inhibits its microtubule-binding activity. JNK3 itself is activated by a series of upstream kinases, as part of a signaling pathway called a MAPK cascade. Our laboratory analyzed changes in the activity of these kinases in the presence of mutant proteins, as well as the resulting phosphorylation in the described signaling pathway.

Funding from NIH (STB), HHMI REU (Hunter College)

Molecular identification of neuron types in the *Xenopus Laevis* vocal circuit

Claire P. Everett[1,3], Darcy B. Kelley[2,3]
[1]Barnard College, New York, NY; [2]Columbia University, New York, NY; [3]Marine Biological Laboratory, Woods Hole, MA.

Patterned behaviors-such as breathing, speaking, and walking- are incorporated seamlessly into everyday life. Diverse neural circuits, endocrine signaling and muscle effectors govern these patterned behaviors. The Kelley lab uses the sexually differentiated vocal behaviors of the South African Clawed frog, *Xenopus laevis*, as a model system to study how male- and female-specific patterns are produced. While some components of vocal circuits have been identified, local hindbrain interneurons and subpopulations of laryngeal motor neurons are poorly understood. Transcription factor expression has been useful in identifying neurons that contribute to locomotion in spinal cord and respiration in the hindbrain. We are using transcription factor and calcium binding protein expression - combined with dye tracing - to identify participants in vocal patterning in *Xenopus*. Calbindin, Isl1, En1 and androgen receptor expression is mapped in late stage tadpole brains using immunohistochemistry (IHC). Specific neurons are identified in the vocal circuit by injecting a fluorescent tracer (rhodaminated dextran) into the vocal motor nucleus and the pattern-generating nucleus and following axon projections and synaptic contacts. Dye-

Nikita , , Han-A , Nelli , , Elizabeth

; ; ;

Alzheimer's Disease (AD) is a neurological disorder characterized by cognitive and motor deterioration that affects memory, attention, speech, and behavior. One of the factors underlying the progression of AD is the accumulation of amyloid beta (Aβ) protein plaques that may block synaptic transmission. It has been previously suggested that mitochondrial dysfunction may play a role in the onset or progression of AD. We now demonstrate that Aβ contributes to cellular dysfunction by affecting both mitochondrial membrane potential and calcium retention capacity -- two measures of mitochondrial function and efficiency. After isolating and purifying mitochondria through density-gradient centrifugation, we found that Aβ-treated mitochondria exhibited a reduction in calcium uptake ability in comparison to control mitochondria. Using confocal microscopy of the mitochondrial membrane potential indicator, TMRE, we visualized and quantified a significant difference in mitochondrial membrane potential between control and Aβ-treated hippocampal neurons. In addition to drastically reducing the membrane potential, Aβ peptide also affected cell morphology – the neurons and mitochondria were irregular in shape, size, frequency, and distribution. We predict that these findings are the result of an Aβ-induced inner mitochondrial membrane leak, and hypothesize that Aβ's interaction with B-cell lymphoma extra-large (Bcl-xL), a specialized Bcl-2 family protein, may cause the opening of the mitochondrial permeability transition pore (mPTP), resulting in decreased mitochondrial competency. Bcl-xL's binding to the beta subunit of the F1F0-ATP synthase reduces the inner membrane ion leak, thereby increasing the flow of ions through the H+ translocator of ATP synthase to make ATP and enhancing the efficiency of energy metabolism in the mitochondria. Since Bcl-xL, among other regulatory proteins, is vital for preventing the neurotoxicity that leads to AD, our findings suggest that Aβ may disrupt regulation of the gating of the inner membrane leak channel within ATP synthase, leading to functional decline of the mitochondria.

Funding by the University of Chicago Jeff Metcalf SURF scholarship

Identifying autotrophic microbes at deep-sea hydrothermal vents using RNA-SIP and RT-qPCR

Paula Pelayo[1], Caroline S. Fortunato[2], Julie A. Huber[2]
[1]UCLA, Los Angeles, CA, [2]Marine Biological Laboratory, Woods Hole, MA.

Deep-sea hydrothermal vents are windows into the microbial community present within the rocky subseafloor of oceanic crust. Subseafloor

Exploring the light environment for larval culturing of MBL squid

Kirsten Peramba[1,2], Scott Bennett

Comparing Scanning Electron Microscopy and Micro-CT Imaging Techniques for the Mechanosensory Lateral Line

Loranzie S. Rogers[1,2], Beth Giuffrida[2,3], Veronique Le Roux[4], Allen F. Mensinger[1,2]
[1]University of Minnesota Duluth, Duluth, MN; [2] Marine Biological Laboratory, Woods Hole, MA, [3]Wareham Middle School, Wareham, MA, [4]Woods Hole Oceanographic Institute, Woods Hole, MA

The lateral line system in fish consists of mechanosensory neuromasts, which detect changes in pressure and vibrations through the water. In this study, scanning electron microscopy (SEM) and micro-CT scanning were conducted to demonstrate how different techniques can be used to visualize and quantify neuromasts along the anterior lateral line (ALL) of Bighead Carp (*Hypophthalmichthys nobilis*), Silver Carp (*H. Molitrix*) and Oyster Toadfish (*Opsanus tau)*. SEM was conducted on Bighead and Silver Carp to map the spatial distribution of neuromasts along the ALL and compare the abundance of superficial neuromasts between the two morphotypes. This technique proved useful in obtaining fine resolution images of hair cells within the neuromasts however was limited to surface analysis only. Micro-CT scanning is a nondestructive 3-D x-ray imaging technique that provides an alternative method to analyze small scale features with less preparation than SEM. Micro-CT scanning was compared using two different micro-CT scanners of varying resolutions which obtained images at pixel sizes of 35.3 and 12.9 m. Scans were then used to create a 3-D model of the anterior portion of the Oyster Toadfish which included the lateral line. This approach allowed for detailed 3-D imaging, but was unable to visualize hair cells with the same detail as SEM. Each technique proved useful at mapping neuromasts, but with structural and detail limitations. The use of these two techniques together allows for hair cell identification and three dimensional reconstruction of neuromasts and the lateral line.

Funding by National Science Foundation Grants IOS-1354745 and DBI-1359230

Using Lipid Biomarkers to Understand Deep Ocean Organic Particle Flux

Leonard Shaw[1], Emily Maness[2], J.C. Weber[3], Maureen Conte[3]
[1]*University of Chicago, Chicago IL;* [2]

Microfluidic Droplets as Substrates for Custom Photolithographic Patterning

Michael J. Sloyan[1,5], Taylor Sulerud[2,5], Ben Noren[2,5], James Pelletier[3,5], Jesse Gatlin[2,5], Christine Field[4,5], Timothy Mitchinson[4,5], John Oakey[2,5]

[1]University of Chicago, Chicago, IL; [2]University of Wyoming, Laramie, WY; [3]Massachusetts Institute of Technology, Cambridge, MA; [4]Harvard Medical School, Boston, MA; [5]Marine Biological Laboratory, Woods Hole, MA

Microfluidic flow devices and positive photolithography are used to create micro-sized structures in polyethylene glycol (PEG) polymers, a biologically inert medium suitable for many scientific and medical uses. The shapes of microfluidic objects, notably droplets, are limited by the internal properties of fluids, whose cohesion encourages simple shapes with radial symmetry. Photolithographic techniques are prized for the intricate patterns they can imprint upon flat films of substrate. Little research has

Tracking biofilm growth of early Plastisphere colonizers

David Vishny[1], Breaun Meeks[2], Erik Zettler[3], Linda Amaral-Zettler[4,5]
[1]*University of Chicago, Chicago, IL,* [2]*Bowie State University, Bowie, MD,* [3]*Sea Education Association, Woods Hole, MA,*
[4]*Marine Biological Laboratory, Woods Hole, MA,*
[5]*Brown University, Providence, RI*

Plastic marine debris (PMD) presents a serious and worsening global pollution problem. PMD permeates even the most distant marine habitats, including deep-sea canyons and polar sea ice layers (Schlining, K, et al., 2013; Obbard, R.W., et al., 2014). Everywhere PMD travels, it comes into contact with microbial life. The surface of PMD provides an exploitable niche for a community dominated by microbes called –the Plastisphere.‖ (Zettler, Mincer and Amaral-Zettler, 2013). While some impacts of PMD on marine animals as consequences of ingestion and entanglement are well-documented, a great number of PMD's ecological impacts remain quite unknown. Understanding the ecology and physiology of the Plastisphere will likely be crucial for our understanding of these so-far-unknown impacts. For example, the behavior of the Plastisphere community may facilitate the transport of invasive algae or bacteria and affect the bioavailability of persistent organic pollutants that adsorb to PMD's hydrophobic surface. Most of the Plastisphere work done so far has been on field samples, which could not be accurately aged. As a result, we are not sure at what rates Plastisphere biofilms form, or how Plastisphere communities change over time, particularly during early community assembly that occurs in the first hours and days of exposure. Our experiments employed lab-grown Plastisphere communities to study biofilm growth rates and interactions between selected microbes on different resins. We found that a non-axenic pennate diatom biofilm formed on four plastic resins at rates greater than one doubling per day, and we found significant differences in growth on different resins. We also found that certain species of Plastisphere bacteria appear to facilitate or inhibit each other's growth in a biofilm.

References

Schlining, K., et al., *Debris in the deep: Using a 22-year video annotation database to survey marine litter in Monterey Canyon, central California, USA.* Deep-Sea Rese416.59 605.79 0 1 36(nsPn161hJET EMC /BT/F5 11.0se4197 505.639m[[(-)] T